

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

FUT3, 35-361aa, Human, His tag, E.coli

产品货号: TP02164

第三版

别名:Fucosyltransferase 3 , CD174, FT3B, FucT-III, LE, Les

描述:Fucosyltransferase 3, also known as FUT3, may catalyze alpha-1,3 and alpha-1,4 glycosidic linkages involved in the expression of Vim-2, Lewis A, Lewis B, sialyl Lewis X and Lewis X/SSEA-1 antigens. This protein may be involved in blood group Lewis determination; Lewis-positive (Le+) individuals have an active enzyme while Lewis-negative (Le-) individuals have an inactive enzyme. Also acts on the corresponding 1,4-galactosyl derivative, forming 1,3-L-fucosyl links. Recombinant human FUT3 protein, fused to His-tag at N-terminus, was expressed in E.coli.

配方:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M urea, 10% glycerol

分子量:40.6 kDa (350aa)

序列:

MGSSHHHHHSSGLVPRGSHMGSRVSRDDATGSPRAPSGSSRQDTPTRPTLLILLWTWPFHIPVALSRCSEMVPGTADC
HITADRKVYPQADTVIVHHWDIMSNPKSRLPPSPRPQGQRWIWFnLEPPNCQHLEALDRYFNLTMSYRSDSDIFTYPYGW
LEPWSGQPAHPPLNLSAKTELVAWAVSNWKPDSARVRYQSLQAHLKVDVYGRSHKPLPKGTMMETLSRYKFYLA FENS
LHPDYITEKLWRNALEAWAVPVVLGPSRSNYERFLPPDAFIHVDDFQSPKDLARYLQELDKDHARYLSYFRWRETLRPRSF
SWALDFCKACWKLQQESRYQTVRSIAAWFT

纯度:> 95% by HPLC

浓度:1 mg/ml (determined by Bradford assay)

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储:+4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.