

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

GREM1, 25-184aa Human, His tag, E.coli

产品货号: TP02332

第三版

别名: Gremlin-1 isoform 1, CKTSF1B1, DAND2, DRM, GREMLIN, IHG-2, PIG2

描述: GREM1 is a member of the BMP (bone morphogenic protein) antagonist family. Like BMPs, BMP antagonists contain cystine knots and typically form homo- and heterodimers. The CAN (cerberus and dan) subfamily of BMP antagonists, to which this protein belongs, is characterized by a C-terminal cystine knot with an eight-membered ring. The antagonistic effect of the secreted glycosylated protein is likely due to its direct binding to BMP proteins. As an antagonist of BMP, this protein may play a role in regulating organogenesis, body patterning, and tissue differentiation. In mouse, this protein has been shown to relay the sonic hedgehog (SHH) signal from the polarizing region to the apical ectodermal ridge during limb bud outgrowth. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. Recombinant human GREM1 protein, fused to His-tag at N-terminus, was expressed in E.coli.

配方: Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

分子量: 20.7 kDa (183aa)

序列:

MGSSHHHHHSSGLVPRGSHMGSKKKGSGAIPPPDKAQHNDSEQTQSPQQPGSRNRGRGQGRGTAMPGEEVLESSQE
ALHVTERKYLKRDWCKTQPLKQTIHEEGCNSRTIINRFCYGQCNSFYIPRHIRKEEGSFQSCSFCKPKKFTTMMVTLNCP
QPPTKKKRVTRVKQCRCISIDL

纯度: > 95% by HPLC

浓度: 0.5 mg/ml (determined by Bradford assay)

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.