

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

HCV-E2, 482-671aa, HCV, His tag, E.coli

产品货号: TP02417

第三版

别名: Polyprotein,

**描述:** E1 and E2 glycoproteins form a heterodimer that is involved in virus attachment to the host cell, virion internalization through clathrin-dependent endocytosis and fusion with host membrane. E1/E2 heterodimer binds to human LDLR, CD81 and SCARB1/SR-BI receptors, but this binding is not sufficient for infection, some additional liver specific cofactors may be needed. The fusion function may possibly be carried by E1. E2 inhibits human EIF2AK2/PKR activation, preventing the establishment of an antiviral state. E2 is a viral ligand for CD209/DC-SIGN and CLEC4M/DC-SIGNR, which are respectively found on dendritic cells (DCs), and on liver sinusoidal endothelial cells and macrophage-like cells of lymph node sinuses. These interactions allow capture of circulating HCV particles by these cells and subsequent transmission to permissive cells. Recombinant HCV (Hepatitis C Virus) E2 protein, fused to His-tag at N-terminus, was expressed in E.coli.

**配方:** Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

**分子量:** 25.4 kDa (226aa)

**序列:**

MRGSHHHHHHGMASMTGGQQMGRDLYDDDDKDRWGSERPWCWHYPPRPGIVPAKSVCGPVCFTSPVVVGTDD  
RSGAPTYSWGANDTDVFLNNTRPPLGNWFGCTWMNSTGFTKVCGAPPCVIGGVGNNTLLCPTDCFRKHPEATYSRC  
GSGPWITPRCMVDYPYRLWHYPCTINYTIFKVRMYVGGVEHRLEAACNWTRGERCDLEDRDRSELSPLLLSTTQ

**纯度:** > 95% by HPLC

**浓度:** 0.25 mg/ml (determined by Bradford assay)

**内毒素:** <1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.