



本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

POLR2C, 1-275aa, Human, His tag, E.coli

产品货号: TP03458

第三版

别名:DNA-directed RNA polymerase II subunit RPB3, hRPB33, hsRPB3, RPB3, RPB31

描述:POLR2C is the third largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. The product of this gene contains a cysteine rich region and exists as a heterodimer with another polymerase subunit, POLR2J. These two subunits form a core subassembly unit of the polymerase. A pseudogene has been identified on chromosome 21. Recombinant human POLR2C protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

配方:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol 1mM DTT, 0.1M NaCl

分子量:33.8 kDa(298aa) confirmed by MALDI-TOF

序列:

MGSSHHHHHSSGLVPRGSHMGSM~~PYANQPTVRITELTDENVKFIIENTDLAVANSIRRVFIAEVPIIAIDWVQIDANSSV~~  
LHDEFIAHRLGLIPLISDDIVDKLQYSR~~DCTCEEFCPECSVEFTLDVRCNEDQTRHVTSDLI~~NSPRV~~IPVTSRNRDNDPND~~  
YVEQDDILIVKLRKGQELRLRAYAKKGFGKEHAKWNPTAGVAFYDPDNALRHTVYPKPEEWPKSEYSELDEDESQAPYD  
PNGKPERFYVNVESCGSLRPETIVLSALSGLKKLSDLQTQLSHEIQSDVLTIN

纯度:> 95% by HPLC

浓度:0.5 mg/ml (determined by Bradford assay)

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.