

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

SELPLG, 42-121aa, Human, His tag, E.coli

产品货号: TP03882

第三版

别名:P-selectin glycoprotein ligand 1 isoform 2 precursor, CD162, CLA, PSGL-1, PSGL1

**描述:**SELPLG is a glycoprotein that functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L-selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response. Alternate splicing results in multiple transcript variants. Recombinant human SELPLG protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

**配方:**Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 10% glycerol, 1mM DTT.

**分子量:**10.9 kDa (103aa) confirmed by MALDI-TOF (Molecular size on SDS-PAGE will appear higher)

**序列:**

MGSSHHHHHSSGLVPRGSHMGSQATEYEYLDYDFLPETEPPEMLRNSTDTTPLTGPGTPESTTVEPAARRSTGLDAGGA  
VTELTTELANMGNLSTDSAAMEI

**纯度:**> 95% by HPLC

**浓度:**0.25 mg/ml (determined by Bradford assay)

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.