

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

STK17A, 64- 321aa Human, His tag, E.coli

产品货号: TP04064

第三版

别名:serine/threonine kinase 17A, DRAK1, DAP kinase-related apoptosis-inducing protein kinase 1, Serine threonine-protein kinase 17A

描述:STK17A is a member of the DAP kinase-related apoptosis-inducing protein kinase family and encodes an autophosphorylated nuclear protein with a protein kinase domain. The protein has apoptosis-inducing activity. Recombinant human STK17A protein, fused to His-tag at N-terminus, was expressed in E.coli.

配方:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 10% glycerol, 0.4M Urea

分子量:31.9kDa (281aa)

序列:

MGSSHHHHHHSSGLVPRGSHMGSGRELGRGKFAVVVRKCIKKDSGKEFAAKFMRKRRKGQDCRMEIHEIAVLELAQDN
PWVINLHEVYETASEMILVLEYAAGGEIFDQC VADREEAFKEKDVQRLMRQILEGVHFLHTRD VVHLDLKPQNILLTSESP
LGDIKIVDFGLSRILKNSEELREIMGTPEYVAPEILSYDPISMATDMWSIGVLTYVMLTGISPFLGNDKQETFLNISQMNL SYS
EEEFDVLSESAVD FIRTLLVKKPEDRATAEECLKHPWL

纯度:> 95% by HPLC

浓度:0.5mg/ml (determined by Bradford assay)

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储:+4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.