

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Activation B7-1 antigen Protein

产品货号: TP04588

第三版

别名:CD80; Activation B7-1 antigen; B7; BB1; CD28LG1; CD28LGB7-1 antigen; T-lymphocyte activation antigen CD80

描述:Cluster of Differentiation 80, also called B7-1, is a member of cell surface immunoglobulin superfamily which plays key, yet distinct roles in the activation of T cells. It is the ligand for two different proteins on the T cell surface: CD28 and CTLA-4. Studies have shown that CTLA-4 binds mostly to CD80. The structure presents two extracellular domains: a membrane distal variable-like domain (IgV) and a membrane proximal Ig constant-like domain (IgC) along with an intracellular domain. Both IgV and IgC consist of anti-parallel beta sandwiches joined by a short linker region. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells.

配方:PBS

分子量:24.7 kDa

序列:Val35-Asn242

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.