

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human EpCAM/TROP1/CD326 Protein(C-Fc)

产品货号: TP04832

第三版

别名: Epithelial Cell Adhesion Molecule; Ep-CAM; Adenocarcinoma-Associated Antigen; Cell Surface Glycoprotein Trop-1; Epithelial Cell Surface Antigen; Epithelial Glycoprotein 314; EGP314; Major Gastrointestinal Tumor-Associated Protein GA733-2; Tumor-Associated Calcium Signal Transducer 1; CD326; EPCAM; GA733-2; TROP1

描述: Epithelial Cell Adhesion Molecule (EpCAM) is a signal type I transmembrane glycoprotein that belongs to the EPCAM family. EpCAM is composed of an extracellular domain with one thyroglobulin type-1 domain, a transmembrane domain and a cytoplasmic domain. EpCAM is found on the surface of adenocarcinoma, but not on mesodermal or neural cell membranes. The EpCAM molecule has been shown to function as a homophilic Ca^{2+} independent adhesion molecule. It may act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium as an immunological barrier providing the first line of defense against infection. Defects in EPCAM are a cause of hereditary non-polyposis colorectal cancer type 8 (HNPCC8) and diarrhea type 5 (DIAR5). EpCAM plays a role in embryonic stem cells proliferation and differentiation; it up-regulates the expression of FABP5, MYC and Cyclin A and Cyclin E. It is highly and selectively expressed by undifferentiated embryonic stem cells.

配方: PBS

分子量: 54.5 kDa

序列: Gln24-Lys265

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.