

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Mouse Activated Leukocyte Cell Adhesion Molecule Protein

产品货号: TP04841

第三版

**别名:**CD166 antigen;cluster of differentiation 166; CD166; activated leucocyte cell adhesion molecule; CD6 ligand; Protein DM-GRASP;CD6L;MEMD

**描述:**Activated leukocyte cell adhesion molecule (ALCAM), also named as CD166 and MEMD, is a type I transmembrane glycoprotein of immunoglobulin superfamily, which mediates homotypic and heterotypic interactions between cells. ALCAM interacts with high affinity with CD6 molecule but weaker homotypic (ALCAM – ALCAM) interactions have also been described. ALCAM – CD6 interactions play an important role in the maintenance of T cell activation, proliferation as well as in formation of immune synapse between antigen-presenting cell and lymphocytes. ALCAM is expressed on a wide variety of cells, particularly on activated lymphocytes, dendritic cells and monocytes, and on various epithelial cell types. It is also involved in multiple processes including embryogenesis, hematopoiesis, angiogenesis, and immune response. While expressed in a wide variety of tissues, ALCAM is usually restricted to subsets of cells in most adult tissues. Recently studies showed ALCAM has prognostic relevance in several human carcinomas, and it has been used as a biomarker for several tumor entities, including melanoma, gynecologic, urologic, and gastrointestinal cancers.

**配方:**PBS

**分子量:**83.3 kDa

**序列:**Met 1-Lys 527

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:**+4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.