

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Triggering Receptor Expressed On Myeloid 2/TREM-2 Protein(C-6His)

产品货号: TP04928

第三版

别名: Triggering receptor expressed on myeloid cells 2; TREM-2; Triggering receptor expressed on monocytes 2; TREM2

描述: Triggering Receptor Expressed on Myeloid cells 2 (TREM2) is a 35 kDa type I transmembrane member of the TREM family and Ig superfamily. Mature human TREM2 consists of a 156 amino acid (aa) extracellular domain (ECD) with one V-type Ig-like domain, a 21 aa transmembrane (TM) domain, and a 35 aa cytoplasmic tail. Soluble forms of the TREM2 ECD are generated by alternative splicing or proteolytic cleavage, and the cytoplasmic domain can be liberated by gamma-Secretase mediated intramembrane cleavage. A positively charged lysine within the transmembrane segment allows association with the signal adapter protein, DAP12 and inhibition of macrophage activation. TREM2 is expressed on macrophages, immature myeloid dendritic cells, osteoclasts, microglia, and adipocytes. It promotes the differentiation and function of osteoclasts, the production of inflammatory cytokines by adipocytes, insulin resistance, and the phagocytic clearance of bacteria.

配方: PBS

分子量: 18.3 kDa

序列: His19-Ser174

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.