

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Rat T-lymphocyte Activation Antigen CD80/B7-1 Protein(C-6His)

产品货号: TP05011

第三版

别名:T-lymphocyte activation antigen CD80; Activation B7-1 antigen; B7; CD80

**描述:**Cluster of Differentiation 80, also called B7-1, is a member of cell surface immunoglobulin superfamily which plays key, yet distinct roles in the activation of T cells. B7-1/CD80 and B7-2/CD86, together with their receptors CD28 and CTLA4, constitute one of the dominant co-stimulatory pathways that regulate T- and B- cell responses. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. Although both CTLA-4 and CD28 can bind to the same ligands, CTLA-4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28 and is involved in the down-regulation of the immune response.

**配方:**PBS

**分子量:**25.7 kDa

**序列:**Vla30-Gln248

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:**+4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.