

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human IL-20 receptor subunit beta/IL-20RB Protein(C-Fc)

产品货号: TP05051

第三版

别名: Interleukin-20 receptor subunit beta; IL-20 receptor subunit beta; IL-20R-beta; IL-20RB; IL-20R2; DIRS1; hCG\_2022374; FNDC6; MGC34923; fibronectin type III domain containing 6; interleukin-20 receptor II

**描述:** Interleukin-20 receptor subunit beta ( IL20RB ) is a single-pass type I membrane protein and belongs to the type II cytokine receptor family. It contains 2 fibronectin type-III domains. There are two kinds of type II cytokine receptors : cytokine receptors that bind type I and type II interferons; cytokine receptors that bind members of the interleukin-10 family (interleukin-10, interleukin-20 and interleukin-22). Type II cytokine receptors are similar to type I cytokine receptors except they do not possess the signature sequence WSXWS that is characteristic of type I receptors. They are expressed on the surface of certain cells, which bind and respond to a select group of cytokines. These receptors are related predominantly by sequence similarities in their extracellular portions that are composed of tandem Ig-like domains. The intracellular domain of type II cytokine receptors is typically associated with a tyrosine kinase belonging to the Janus kinase (JAK) family

**配方:** PBS

**分子量:** 49.6 kDa

**序列:** Asp30-Ala230

**纯度:** > 95% by HPLC

**浓度:**

**内毒素:** <1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.