

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Interleukin-17F/IL-17F Protein(C-6His)

产品货号: TP05366

第三版

别名: Interleukin-17F; IL-17F; Cytokine ML-1; Interleukin-24; IL-24; IL17F; IL24

描述: Interleukin-17F (IL-17F) exists in a disulfide-linked heterodimer that belongs to the IL-17 family. IL-17F is expressed in activated, but not resting, CD4+ T-cells and activated monocytes. IL-17F has been shown to stimulate the production of several other cytokines, including IL-6, IL-8, and granulocyte colony-stimulating factor. IL-17F can regulate cartilage matrix turnover and stimulates PBMC and T-cell proliferation. IL-17F is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. Defects in IL-17F are the cause of familial candidiasis type 6 (CANDF6).

配方: PBS

分子量: 16.0 kDa

序列: Arg31-Gln163

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.