

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Epidermal Growth Factor Receptor Protein

产品货号: TP05436

第三版

别名: Epidermal growth factor receptor; EGFR; Proto-oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1; EGFR

描述: The EGFR subfamily of receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43% - 44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glycoproteins with an extracellular ligand binding domain. The extracellular ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF- α , betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.

配方: PBS

分子量: 65.8 kDa

序列: Leu25-Val30Gly&Asn298-Ser645

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 μ g of protein (determined by LAL method)

存储: +4 °C 保存 (1-2 周). 长期保存在 -20 °C 或者 -70 °C. 避免反复冻融.