

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human C-C Motif Chemokine 16/CCL16 Protein

产品货号: TP05568

第三版

别名:C-C Motif Chemokine 16; Chemokine CC-4; HCC-4; Chemokine LEC; IL-10-Inducible Chemokine; LCC-1; Liver-Expressed Chemokine; Lymphocyte and Monocyte Chemoattractant; LMC; Monotactin-1; MTN-1; NCC-4; Small-Inducible Cytokine A16; CCL16; ILINCK; NCC4; SCYA16

描述:CCL16 is a member of CC chemokine family. CCL16 cDNA encodes a 120 amino acid peptide along with a 23 amino acids signal peptide that is cleaved to generate 97 amino acid protein. CCL16 is distantly related to other CC chemokines, showing less than 30% sequence identity. CCL16 elicits its effects on cells by interacting with cell surface chemokine receptors such as CCR1, CCR2, CCR5 and CCR8. Recombinant CCL16 has been shown to chemoattract human monocytes and THP1 cells but not resting lymphocytes nor neutrophils. CCL16 has potent myelosuppressive activity, suppresses proliferation of myeloid progenitor cells. CCL16 induces a calcium flux in THP1 cells that can be desensitized by prior exposure to RANTES, suggesting that CCL16 and RANTES share the same receptor in THP1 cells.

配方:PBS

分子量:11.0 kDa

序列:Gln24-Gln120

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.