
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Mouse FLT-3 / CD135 / FLK-2 Protein (His tag)

产品货号: TP05604

第三版

别名: B230315G04; CD135; Fik-2; Fik2; Flt-3; Ly72; wmf1

描述: The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD135, also known as FLT-3, FLK-2, is a member of the CD system. CD135 is an important cell surface marker recognized by specific sets of antibodies to identify the types of hematopoietic (blood) progenitors in the bone marrow and its function to differentiate hematopoietic stem cells, which are CD135 negative, from multipotent progenitors, which are CD135 positive. CD135 is a receptor tyrosine kinase type α ϕ for the cytokine Flt3 ligand and activates signaling through second messengers by binding to Flt3. Signaling through CD135 is important for lymphocyte development. The encoding gene CD135 is a proto-oncogene to which mutations happened will lead to cancer such as leukemia. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

配方: PBS

分子量: 59.5 kDa

序列: Met1-Ser544

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.