

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Isocitrate Dehydrogenase 1/IDH1 Protein(C-8His)

产品货号: TP05960

第三版

别名:Isocitrate Dehydrogenase [NADP] Cytoplasmic; IDH; Cytosolic NADP-Isocitrate Dehydrogenase; IDP; NADP(+)-Specific ICDH; Oxalosuccinate Decarboxylase; IDH1; PICD

描述:Isocitrate Dehydrogenase [NADP] Cytoplasmic (IDH1) belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 exists as a homodimer, binding one magnesium or manganese ion per subunit. Mutations of IDH1 have been shown to cause metaphyseal chondromatosis with aciduria and are involved in the development of glioma IDH plays a role in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the α -hydroxylation of phytanic acid.

配方:PBS

分子量:48.1 kDa

序列:Met 1-Leu414

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.