

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human T cell Immunoreceptor with Ig and ITIM domains Protein

产品货号: TP06048

第三版

**别名:**T-cell immunoreceptor with Ig and ITIM domains;;VSI9; VSTM3;TIGIT;V-set and transmembrane domain-containing protein 3;V-set and immunoglobulin domain-containing protein 9

**描述:**T cell immunoreceptor with Ig and ITIM domains (TIGIT) is a member of the CD28 family within the Ig superfamily of proteins. TIGIT is expressed on NK cells and subsets of activated, memory and regulatory T cells, and particularly on follicular helper T cells within secondary lymphoid organs. It binds to CD155 and Nectin-2 that appear on dendritic cells (DC) and endothelium. Ligation of TIGIT on T cells down-regulates TCR-mediated activation and subsequent proliferation, while NK cell TIGIT ligation blocks NK cell cytotoxicity. Through CD155 and Nectin-2, which also interact with DNAM-1/CD226 and CD96/Tactile, TIGIT is part of an interacting network of Ig superfamily members that may augment or oppose each other. In particular, TIGIT binding to CD155 can antagonize the effects of DNAM1.

**配方:**PBS

**分子量:**14.1 kDa

**序列:**Met22-Pro141

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.