

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant HumanCD40L/CD154/TNFSF5/protein(His Tag)

产品货号: TP06213

第三版

别名:CD154;CD40 Ligand;CD40L;gp39;hCD40L;HIGM1;IGM;IMD3;T-BAM;TNFSF5;TRAP

描述:The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD154, also known as CD40 ligand or CD40L, is a member of the TNF superfamily. While CD154 was originally found on T cell surface, its expression has since been found on a wide variety of cells, including platelets, mast cells, macrophages and NK cells. CD154's ability is achieved through binding to the CD40 on antigen- presenting cells (APC). In the macrophage cells, the primary signal for activation is IFN- γ from Th1 type CD4 T cells. The secondary signal is CD40L on the T cell, which interacting with the CD40 molecules, helping increase the level of activation. Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Proteins Immune Checkpoint Targets Co-stimulatory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy

配方:PBS

分子量:17.7 kDa

序列:Glu 108-Leu 261

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.