

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human CLEC4A / CLECSF6 / DCIR Protein (His tag)

产品货号: TP06534

第三版

别名:CD367;CLEC4A;CLECSF6;DCIR;DDB27;HDCGC13P;LLIR

描述:Dendritic cell immunoreceptor (DCIR), also known as C-type lectin domain family 4 member A (CLEC4A), C-type lectin superfamily member 6 (CLECSF6), is a single-pass type II C-type lectin receptor expressed mainly in dendritic cells (DCs), which is a negative regulator of DC expansion and has a crucial role in maintaining the homeostasis of the immune system. The Dectin-2 family of C-type lectins that includes Dectin-2, BDCA-2, DCIR, DCAR, Clecsf8 and Mincle. These type II receptors contain a single extracellular carbohydrate recognition domain and have diverse functions in both immunity and homeostasis. DCIR is the only member of the family which contains a cytoplasmic signalling motif and has been shown to act as an inhibitory receptor, while BDCA-2, Dectin-2, DCAR and Mincle all associate with FcRgamma chain to induce cellular activation, including phagocytosis and cytokine production. Dectin-2 and Mincle have been shown to act as pattern recognition receptors for fungi, while DCIR acts as an attachment factor for HIV. In addition to pathogen recognition, DCIR has been shown to be pivotal in preventing autoimmune disease by controlling dendritic cell proliferation. DCIR expressed on antigen presenting cells and granulocytes and acts as an inhibitory receptor via an intracellular immunoreceptor tyrosine-based inhibitory motif (ITIM). It may also be involved via its ITIM motif in the inhibition of B-cell-receptor-mediated calcium mobilization and protein tyrosine phosphorylation. Additionally, DCIR can participate in the capture of HIV-1 and promote infection in trans and in cis of autologous CD4(+) T cells from human immature monocyte-derived DCs. DCIR acts as a ligand for HIV-1 and is involved in events leading to productive virus infection.

配方:PBS

分子量:22 kDa

序列:Gln 70-Leu 237

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储:+4 ° C 保存 (1-2 周). 长期保存在-20 ° C 或者-70 ° C. 避免反复冻融.