

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Peroxiredoxin 2 / PRDX2 Protein (His tag)

产品货号: TP06625

第三版

别名:HEL-S-2a;NKEF-B;NKEFB;PRP;PRX2;PRXII;PTX1;TDPX1;TPX1;TSA

描述: Peroxiredoxin-2, also known as Natural killer cell-enhancing factor B, NKEF-B, Thiol-specific antioxidant protein, Thioredoxin peroxidase 1, Thioredoxin-dependent peroxide reductase 1, PRDX2 and NKEFB, is a cytoplasm protein which belongs to the ahpC / TSA family. Peroxiredoxin-2 / PRDX2 contains one thioredoxin domain. Peroxiredoxin-2 / PRDX2 is involved in redox regulation of the cell. It reduces peroxides with reducing equivalents provided through the thioredoxin system. Peroxiredoxin-2 / PRDX2 is not able to receive electrons from glutaredoxin. It may play an important role in eliminating peroxides generated during metabolism. Peroxiredoxin-2 / PRDX2 might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H₂O₂. The Peroxiredoxins / Prx are a family of peroxidases that can reduce H₂O₂ using an electron from thioredoxin (Trx) or other substances. The mammalian Peroxiredoxins / Prx family is divided into six groups (PRDX1 , PRDX2, PRDX3, PRDX4, PRDX5, PRDX6) on the basis of homology of amino acid sequences. They are located in the cytosol and play a role in the cell signaling system. All six mammalian peroxiredoxins are expressed in the lung. Peroxiredoxins / Prx is overexpressed in breast cancer tissues to a great extent suggesting that Peroxiredoxins / Prx has a proliferative effect and may be related to cancer development or progression.

配方: PBS

分子量: 24 kDa

序列: Met 1-Asn 198

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.