

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Rat GP1BB / CD42c Protein (Fc tag)

产品货号: TP06796

第三版

别名:GP1BB

描述:Platelet glycoprotein Ib (GPIb) complex is best known as a major platelet receptor for von Willebrand factor essential for platelet adhesion under high shear conditions found in arteries and in thrombosis. The GPIb complex is composed of GPIb alpha (Platelet glycoprotein Ib alpha chain) covalently attached to GPIb beta (Platelet glycoprotein Ib beta chain) and noncovalently complexed with GPIX and GPV. GPIb-beta, also known as GP1BB, CD42b-beta and CD42c, is single-pass type I membrane protein expressed in heart and brain, which is a critical component of the von Willebrand factor (vWF) receptor. The cysteine knot region of GPIb beta in the N terminus is critical for the conformation of GPIb beta that interacts with GPIX. The precursor of GP1BB is synthesized from a 1.0 kb mRNA expressed in platelets and megakaryocytes. GPIb is a heterodimeric transmembrane protein consisting of a disulfide-linked 140 kD alpha chain and 22 kD beta chain. GPIb alpha chain provides the vWF binding site, and GPIb beta chain contributes to surface expression of the receptor and participates in transmembrane signaling through phosphorylation of its intracellular domain. GP1BB is part of the GPIb-V-IX system that constitutes the receptor for von Willebrand factor (vWF), and mediates platelet adhesion in the arterial circulation. Defects in GP1BB are a cause of Bernard-Soulier syndrome (BSS), also known as giant platelet disease (GPD). BSS patients have unusually large platelets and have a clinical bleeding tendency.

配方:PBS

分子量:40.1 kDa

序列:Met1-Cys147

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.