

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human IκB alpha / NFKBIA Protein (His tag)

产品货号: TP06854

第三版

别名:IKBA;MAD-3;NFKBI

**描述:**Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (IκB alpha, NFKBIA, or IKBA), is a member of the NF-kappa-B inhibitor family that function to inhibit the NF-kB transcription factor. NFKBIA inhibits NF-kB by masking the nuclear localization signals (NLS) of NF-kB proteins and keeping them sequestered in an inactive state in the cytoplasm. In addition, NFKBIA blocks the ability of NF-κB transcription factors to bind to DNA, which is required for NF-kB's proper functioning. Signal-induced degradation of I kappa B alpha exposes the nuclear localization signal of NF-kappa B, thus allowing it to translocate into the nucleus and activate transcription from responsive genes. An autoregulatory loop is established when NF-kappa B induces expression of the I kappa B alpha gene and newly synthesized I kappa B alpha accumulates in the nucleus where it negatively regulates NF-kappa B-dependent transcription. As part of this post-induction repression, the nuclear export signal on I kappa B alpha mediates transport of NF-kappa B-I kappa B alpha complexes from the nucleus to the cytoplasm. Deletion of NFKBIA has an effect that is similar to the effect of EGFR amplification in the pathogenesis of glioblastoma and is associated with comparatively short survival. Polymorphisms in NFKBIA may be important in pre-disposition to and outcome after treatment, of multiple myeloma (MM). The NFKBIA gene product, IκappaBα, binds to NF-kappaB preventing its activation and is important in mediating resistance to apoptosis in B-cell lymphoproliferative diseases.

**配方:**PBS

**分子量:**36.4 kDa

**序列:**Phe 2-Leu 317

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.