

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human RET Kinase Protein (His tag)

产品货号: TP06875

第三版

别名:CDHF12;CDHR16;HSCR1;MEN2A;MEN2B;MTC1;PTC;RET-ELE1;RET51

描述:RET proto-oncogene, also known as RET, is a cell-surface molecule that transduce signals for cell growth and differentiation. It contains 1 cadherin domain and 1 protein kinase domain. RET proto-oncogene belongs to the protein kinase superfamily, tyrosine protein kinase family. RET proto-oncogene is involved in numerous cellular mechanisms including cell proliferation, neuronal navigation, cell migration, and cell differentiation upon binding with glial cell derived neurotrophic factor family ligands. It phosphorylates PTK2/FAK1 and regulates both cell death/survival balance and positional information. RET is required for the molecular mechanisms orchestration during intestine organogenesis; involved in the development of enteric nervous system and renal organogenesis during embryonic life; promotes the formation of Peyer's patch-like structures; modulates cell adhesion via its cleavage; involved in the development of the neural crest. RET proto-oncogene is active in the absence of ligand, triggering apoptosis. RET acts as a dependence receptor; in the presence of the ligand GDNF in somatotrophs (within pituitary), promotes survival and down regulates growth hormone (GH) production, but triggers apoptosis in absence of GDNF. It also regulates nociceptor survival and size; triggers the differentiation of rapidly adapting (RA) mechanoreceptors; mediated several diseases such as neuroendocrine cancers. Defects in RET may cause colorectal cancer, hirschsprung disease type 1, medullary thyroid carcinoma, multiple neoplasia type 2B, susceptibility to pheochromocytoma, multiple neoplasia type 2A, thyroid papillary carcinoma and congenital central hypoventilation syndrome. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

配方:PBS

分子量:69.1 kDa

序列:Met 1-Arg 635

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储:+4 ° C 保存 (1-2 周). 长期保存在-20 ° C 或者-70 ° C. 避免反复冻融.