

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human IL-8 / CXCL8 Protein (aa 28-99, Fc tag)

产品货号: TP07025

第三版

别名:GCP-1;GCP1;IL-8;IL8;Interleukin-8;LECT;LUCT;LYNAP;MDNCF;MONAP;NAF;NAP-1;NAP1

**描述:**Interleukin 8 (IL-8), also known as CXCL8, which is a chemokine with a defining CXC amino acid motif that was initially characterized for its leukocyte chemotactic activity, is now known to possess tumorigenic and proangiogenic properties as well. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to inflammatory stimuli (IL1, TNF, LPS, etc). In human gliomas, IL-8 is expressed and secreted at high levels both in vitro and in vivo, and recent experiments suggest it is critical to glial tumor neovascularity and progression. Levels of IL-8 correlate with histologic grade in glial neoplasms, and the most malignant form, glioblastoma, shows the highest expression in pseudopalisading cells around necrosis, suggesting that hypoxia/anoxia may stimulate expression. Interleukin (IL)-8/CXCL8 is a potent neutrophil chemotactic factor. Accumulating evidence has demonstrated that various types of cells can produce a large amount of IL-8/CXCL8 in response to a wide variety of stimuli, including proinflammatory cytokines, microbes and their products, and environmental changes such as hypoxia, reperfusion, and hyperoxia. Numerous observations have established IL-8/CXCL8 as a key mediator in neutrophil-mediated acute inflammation due to its potent actions on neutrophils. However, several lines of evidence indicate that IL-8/CXCL8 has a wide range of actions on various types of cells, including lymphocytes, monocytes, endothelial cells, and fibroblasts, besides neutrophils. The discovery of these biological functions suggests that IL-8/CXCL8 has crucial roles in various pathological conditions such as chronic inflammation and cancer. IL-8 has been associated with tumor angiogenesis, metastasis, and poor prognosis in breast cancer. IL-8 may present a novel therapeutic target for estrogen driven breast carcinogenesis and tumor progression.

**配方:**PBS

**分子量:**35 kDa

**序列:**Ser 28-Ser 99

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:**+4 ° C 保存 (1-2 周). 长期保存在-20 ° C 或者-70 ° C. 避免反复冻融.