

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human GAD65 / GAD2 / GAD-2 Protein (GST Tag)

产品货号: TP07040

第三版

别名:GAD65

**描述:**Glutamate decarboxylase 2, also known as glutamate decarboxylase 65 kDa isoform, 65 kDa glutamic acid decarboxylase, GAD2 and GAD65, is a member of the group II decarboxylase family. GAD2 is identified as a major autoantigen in insulin-dependent diabetes. GAD2 is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. GAD2 may also play a role in the stiff man syndrome. GAD2 is implicated in the formation of the gamma-aminobutyric acid (GABA), a neurotransmitter involved in the regulation of food intake. GABA is synthesized in brain by two isoforms of glutamic acid decarboxylase (Gad), GAD1 and GAD2. GAD1 provides most of the GABA in brain, but GAD2 can be rapidly activated in times of high GABA demand. Mice lacking GAD2 are viable whereas deletion of GAD1 is lethal. Deletion of GAD2 increased ethanol palatability and intake and slightly reduced the severity of ethanol-induced withdrawal. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

**配方:**PBS

**分子量:**92.6 kDa

**序列:**Met 1-Leu 585

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.