

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Recombinant Mouse FABP4 / ALBP / A-FABP Protein (His & MYC Tag)

产品货号: TP07051

第三版

别名:422/aP2;ALBP/Ap2;Ap2;Lbpl

描述:Fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, Fatty acid-binding protein, Fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, Fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, Fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, Fatty acid-binding protein, adipocyte, also known as Adipocyte-type fatty acid-binding protein, adipocyte-type Adipocyte lipid-binding protein, and FABP4, is a cytoplasm protein which belongs to the calycin superfamily and Fatty-acid binding protein (FABP) family. In familial combined hyperlipidemia (FCHL), FABP4 correlated to body mass index (BMI), waist circumference and homeostasis model assessment (HOMA) index.FABP4 levels were associated with triglyceride-rich lipoproteins. In humans serum FABP4 levels correlate significantly with features of PCOS. It appears to be a determinant of atherogenic dyslipidemia. FABP4 pathway mediates the sebaceous gland hyperplasia in keratinocyte-specific Pten-null mice. FABP4 concentration significantly increased with an increasing of MS features and was strongly correlated with body-mass index, triglycerides, HDL-cholesterol concentrations and blood pressure. Patients in the highest quartile of FABP4 presented a six-fold increased odds ratio for MS and a three-fold increased odds for LD, adjusted by age, sex, body-mass index and the antiretroviral therapy. FABP4 is a strong plasma marker of metabolic disturbances in HIV-infected patients, and therefore, could serve to guide therapeutic intervention in this group of patients.

配方:PBS

分子量:16.5 kDa

序列:Cvs2-Ala132

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4°C保存(1-2周).长期保存在-20°C或者-70°C.避免反复冻融.

1/1