

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Mouse Epcr / PROCR Protein (Fc tag)

产品货号: TP07118

第三版

别名: AI325044; Ccca; Ccd41; Epcr

描述: Endothelial protein C receptor (EPCR), also known as activated protein C receptor (APC receptor) or PROCR, is a receptor for Protein C. Protein C plays an important role in many metabolism processes in humans and other animals after activated by binding to Endothelial protein C receptor (EPCR). Because of the EPCR is found primarily on endothelial cells (cells on the inside of blood vessels), activated protein C is found mainly near endothelial cells. Protein C is pleiotropic, with two main functions: anticoagulation and cytoprotection. Which function will be performed depends on whether or not protein C remains bound to EPCR after activated. The anticoagulation occurs when it does not. In this case, protein C functions as an anticoagulant by irreversibly proteolytically inactivating Factor Va and Factor VIIIa, turning them into Factor Vi and Factor VIIIi respectively. When still bound to EPCR, activated protein C performs its cytoprotective effects, acting on the effector substrate PAR-1, protease-activated receptor-1. To a degree, APC's anticoagulant properties are independent of its cytoprotective ones, in that expression of one pathway is not affected by the existence of the other.

配方: PBS

分子量: 49.3 kDa

序列: Met 1-Ser 214

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.