
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human TIM4 / TIMD4 Protein (His tag)

产品货号: TP07188

第三版

别名:SMUCKLER;TIM4

描述:A type I transmembrane protein called TIM4 (T-cell immunoglobulin- and mucin-domain-containing molecule; also known as TIMD4), which belongs to the immunoglobulin superfamily and TIM family. TIM4 is involved in regulating T-cell proliferation and lymphotoxin signaling. It is a ligand for HAVCR1/TIMD1. Recent reports indicate that dendritic cell (DC)-derived T-cell immunoglobulin and mucin domain molecule (TIM)-4, which is expressed on dendritic cells and macrophages, plays an important role in the initiation of T(H)2 polarization. TIM4 bound apoptotic cells by recognizing phosphatidylserine via its immunoglobulin domain. The expression of TIM4 in fibroblasts enhanced their ability to engulf apoptotic cells. TIM4 is phosphatidylserine receptor for the engulfment of apoptotic cells, and may also be involved in intercellular signalling in which exosomes are involved. Modulation of TIM4 production in dendritic cells (DCs) represents a novel therapeutic approach for the treatment of peanut allergy. The interaction of TIM1/TIM4 played a critical role in sustaining the polarization status of Th2 cells in allergic rhinitis (AR) patients. Cross-linking FcγRI by antigen/IgG complexes increased the production of TIM4 by dendritic cells via upregulating tumor necrosis factor-α in DCs. Specific immunotherapy (SIT) suppresses the skewed Th2 responses via disrupting the interaction of TIM1/TIM4 in antigen-specific Th2 cells.

配方:PBS

分子量:32.7 kDa

序列:Met 1-Leu 315

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.