
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Rat CD64 / FCGR1A Protein (His tag)

产品货号: TP07189

第三版

别名:FcgammaRI;Fcgr1;Fcgr1b

描述:High affinity immunoglobulin gamma Fc receptor I, also known as FCGR1 and CD64, is an integral membrane glycoprotein and a member of the immunoglobulin superfamily. CD64 is a high affinity receptor for the Fc region of IgG gamma and functions in both innate and adaptive immune responses. Receptors that recognize the Fc portion of IgG function in the regulation of immune response and are divided into three classes designated CD64, CD32, and CD16. CD64 is structurally composed of a signal peptide that allows its transport to the surface of a cell, three extracellular immunoglobulin domains of the C2-type that it uses to bind antibody, a hydrophobic transmembrane domain, and a short cytoplasmic tail. CD64 is constitutively found on only macrophages and monocytes, but treatment of polymorphonuclear leukocytes with cytokines like IFN γ and G-CSF can induce CD64 expression on these cells. The inactivation of the mouse CD64 resulted in a wide range of defects in antibody Fc-dependent functions. Mouse CD64 is an early participant in Fc-dependent cell activation and in the development of immune responses.

配方:PBS

分子量:31.6 kDa

序列:Met 1-Pro 285

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.