

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Recombinant Human SIRP alpha / CD172a Protein (His tag)

产品货号: TP07314

第三版

## 别名:BIT;CD172A;MFR;MYD-1;P84;PTPNS1;SHPS1;SIRP;SIRP alpha

描述:Tyrosine-protein phosphatase non-receptor type substrate 1, also known as SHP substrate 1, Inhibitory receptor SHPS-1, Brain Ig-like molecule with tyrosine-based activation motifs, Macrophage fusion receptor, CD172 antigen-like family member A, SIRPA and CD172a, is a single-pass type I membrane protein which contains two Ig-like C1-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. SIRPA is ubiquitously expressed. It is highly expressed in brain and detected at lower levels in heart, placenta, lung, testis, ovary, colon, liver, small intestine, prostate, spleen, kidney, skeletal muscle and pancreas. It is also detected on myeloid cells, but not T-cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRPA acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. It may play a key role in intracellular signaling during synaptogenesis and in synaptic function. SIRPA is involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. It mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation.Immune Checkpoint Detection: IP Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Targets Co-inhibitory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy

配方:PBS

**分子量**:39 kDa

序列:Met 1-Arg 370

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4°C保存 (1-2周). 长期保存在-20°C或者-70°C. 避免反复冻融.