

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human S100A8 Protein (His Tag)

产品货号: TP07580

第三版

别名:60B8AG;CAGA;CFAG;CGLA;CP-10;L1Ag;MA387;MIF;MRP-8;MRP8;NIF;P8

**描述:**S100A8 is a member of the S100 protein family containing 2EF-hand calcium-binding motifs. S100 proteins are involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. Altered expression of S100A8 protein is associated with various diseases and cancers. S100A8 may have an immunoregulatory role by contributing to the regulation of fetal-maternal interactions. It may play a protective role and its absence may allow infiltration by maternal cells, a process eventually manifesting as resorption. The heterodimeric S100 protein complex S100A8/A9 which has been shown to be involved in inflammatory and neoplastic disorders. The complex can induce cell proliferation, or apoptosis, inflammation, collagen synthesis, and cell migration. S100A8/A9 has emerged as important pro-inflammatory mediator in acute and chronic inflammation. More recently, increased S100A8 and S100A9 levels were also detected in various human cancers, presenting abundant expression in neoplastic tumor cells as well as infiltrating immune cells. On the one hand, S100A8/A9 is a powerful apoptotic agent produced by immune cells, making it a very fascinating tool in the battle against cancer. It spurs the risk to induce auto-immune response and may serve as a lead compound for cancer-selective therapeutics. In contrast, S100A8/A9 expression in cancer cells has also been associated with tumor development, cancer invasion or metastasis. Altogether, its expression and potential cytokine-like function in inflammation and in cancer suggests that S100A8/A9 may play a key role in inflammation-associated cancer.

**配方:**PBS

**分子量:**12.2 kDa

**序列:**Met 1-Glu 93

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.