

---

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

---

Recombinant Human FLRT2 Protein (His Tag)

产品货号: TP07625

第三版

别名:KIAA0405;UNQ232/PRO265

**描述:**Fibronectin Leucine-Rich Transmembrane (FLRT) proteins are glycosylated membrane proteins expressed at the cell surface which localise in a homophilic manner to cell-cell contacts expressing the focal adhesion marker vinculin. FLRT1, FLRT2, and FLRT3, the three genes encode putative type I transmembrane proteins, each containing 10 leucine-rich repeats (LRR), a type III fibronectin (FN) domain, followed by the transmembrane region, and a short cytoplasmic tail. FLRT family members may function in cell adhesion and/or receptor signalling. Each member of the FLRT family has a distinct, highly regulated expression pattern, as was seen for the NLRR family. FLRT2 is expressed in a subset of the sclerotome, adjacent to the region that forms the syndetome, suggesting that interaction with FGF signalling may be a general property of FLRT proteins. All FLRTs can interact with FGFR1 and FLRTs can be induced by the activation of FGF signalling by FGF-2. FLRT proteins have a dual role, promoting FGF signalling and modulating homotypic cell adhesion. FLRT2 played critical roles in craniofacial development, and it was also present in the vomero-nasal organ, mandibular primordia, and the posterior aspects of the unfused and fused secondary palatal shelves.

**配方:**PBS**分子量:**57.7 kDa**序列:**Met 1-Ser 539**纯度:**> 95% by HPLC**浓度:****内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.