

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Recombinant Mouse TRAIL R2 / CD262 / TNFRSF10B Protein (His tag)

产品货号: TP07663

第三版

## 别名:DR5;KILLER;Ly98;MK;TRAILR2;TRICK2A;TRICK2B;TRICKB

描述:Tumor necrosis factor receptor superfamily, member 10b, official symbol TNFRSF10B, also known as Death receptor 5, CD262, TNF-related apoptosis-inducing ligand receptor 2 (TRAIL R2), is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. TRAIL R2/CD262/TNFRSF10B was purified independently as the only receptor for TRAIL detectable on the surface of two different human cell lines that undergo apoptosis upon stimulation with TRAIL. TRAIL R2/CD262/TNFRSF10B contains two extracellular cysteine-rich repeats, typical for TNF receptor (TNFR) family members, and a cytoplasmic death domain. TRAIL R2/CD262/TNFRSF10B mediates apoptosis via the intracellular adaptor molecule FADD/MORT1. TRAIL receptors can signal both death and gene transcription, functions reminiscent of those of TNFR1 and TRAMP, two other members of the death receptor family. Defects in TRAIL R2/CD262/TNFRSF10B may be a cause of head and neck squamous cell carcinomas (HNSCC) also known as squamous cell carcinoma of the head and neck.Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

配方:PBS

**分子量**:15 kDa

序列:Met 1-Ser 177

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4°C保存 (1-2周).长期保存在-20°C或者-70°C.避免反复冻融.