
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human CD89 / FCAR Protein (His tag)

产品货号: TP07873

第三版

别名:CD89;CTB-61M7.2;FcalphaRI;FCAR;XXbac-BPG230H20.5

描述:FCAR, also called Fc RI or CD89, is a type I transmembrane receptor for Fc region of IgA which is the most abundant immunoglobulin in mucosal areas but is only the second most common antibody isotype in serum. This receptor is present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, especially phagocytes located in mucosal areas. Upon ligand IgA binding, Fc RI associates with the FcR signaling molecule bearing the immunoreceptor tyrosine-based activation motif (ITAM) through a unique charge-based mechanism and triggers multiple cell-mediated immune responses. It has been reported that Fc RI is a dual-function receptor that can mediate both inflammatory and anti-inflammatory responses depending on the type of interaction with its ligand. Sustained aggregation of FCAR results in activation of target-cell functions such as antigen presentation and cytokine release. In contrast, Monomeric targeting with serum IgA or with a variety of anti-Fc RI Fab fragments triggers an inhibitory response and additionally induces apoptosis. Fc RI thus play an fundamental role in preventing tumor development and growth, as well as in controlling inflammation.

配方:PBS

分子量:25 kDa

序列:Met 1-Asn 227

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.