

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human FGFR1 / CD331 Protein (His & Fc tag)

产品货号: TP07924

第三版

别名:bFGF-R-1;BFGFR;CD331;CEK;FGFBR;FGFR-1;FLG;FLT-2;FLT2;HBGFR;HH2;HRTFDS;KAL2;N-SAM;OGD

描述:FGFR1, also known as CD331, belongs to the fibroblast growth factor receptor subfamily where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. Fibroblast growth factors (FGFs) (FGF1 - 10 and 16 - 23) are mitogenic signaling molecules that have roles in angiogenesis, wound healing, cell migration, neural outgrowth and embryonic development. FGFs bind heparan sulfate glycosaminoglycans, which facilitates dimerization (activation) of FGF receptors. FGFR1 is a full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR1 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. CD331 can be detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells. Defects in FGFR1 are a cause of Pfeiffer syndrome , idiopathic hypogonadotropic hypogonadism , Kallmann syndrome type 2, osteoglophonic dysplasia and trigonocephaly non-syndromic.

配方:PBS

分子量:57.5 kDa

序列:Met 1-Glu 285

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.