

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human Pepsinogen C / PGC Protein (His tag)

产品货号: TP07991

第三版

别名:PEPC;PGII

**描述:**Pepsinogen C, also known as PGC, is an aspartic proteinase that belongs to the peptidase family A1. Pepsinogen C is synthesized in the gastric mucosa as inactive precursors, known as zymogens. Pepsinogen C contains a prosegment that serves to stabilize the inactive form and prevent entry of the substrate to the active site. At low PH conditions, Pepsinogen C undergoes conversion into active enzyme. Pepsinogen C has been found expressed in all regions of the stomach mucosa and also in the proximal duodenal mucosa. In stomach cancer tissues and cancer cell lines, the expressions of the pepsinogen genes were decreased or lost, in good accordance with their pepsinogen productions. No gross structural changes of the pepsinogen genes were observed in these cancers, but the methylation patterns of the pepsinogen genes were found to be altered in different ways in different cancers. Serum levels of Pepsinogen C are used as a biomarker for certain gastric diseases including Helicobacter pylori related gastritis.

**配方:**PBS

**分子量:**42 kDa

**序列:**Met 1-Ala 388

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.