

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human PAPP2 / Pappalysin 2 Protein (His tag)

产品货号: TP08004

第三版

别名:PAPP-A2;PAPP-E;PAPPE;PLAC3

**描述:**Pappalysin-2/PAPP-A2 is the second member of the pappalysin family of metzincin superfamily, of which PAPP-A is the first member. There is no homology between the prepro-peptides of PAPP-A and PAPP-A2, but 46% of the residues of mature PAPP-A are also present in mature PAPP-A2. PAPP-A specifically cleaves insulin-like growth factor-binding protein(IGFBP)-4, one of six known modulators of IGF-I and -II, whereas PAPP-A2 specifically cleaved IGFBP-5 at one site, between Ser-143 and Lys-144. In contrast to the cleavage of IGFBP-4 by PAPP-A that strictly requires the presence of IGF, the cleavage of IGFBP-5 by PAPP-A2 was IGF-independent. Recent data firmly establish PAPP-A and IGFBP-4 as an important functional pair in several systems. Because of its close relationship with PAPP-A, both structurally and functionally, PAPP-A2 is a likely candidate for IGFBP-5 proteinase in many tissues and conditioned media where IGFBP-5 proteolysis has been reported.

**配方:**PBS

**分子量:**131 kDa

**序列:**Ser 234-Cys 1396

**纯度:**> 95% by HPLC

**浓度:**

**内毒素:**<1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.