

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human KEAP1 / INRF2 Protein

产品货号: TP08086

第三版

别名: INRF2; KEAP-1; KLHL19

描述: Kelch-like ECH-associated protein 1, also known as cytosolic inhibitor of Nrf2, Kelch-like protein 19, KEAP1 and INRF2, is a cytoplasm and nucleus protein which contains one BACK (BTB/Kelch associated) domain, one BTB (POZ) domain and six Kelch repeats. KEAP1 / INRF2 is broadly expressed, with highest levels in skeletal muscle. KEAP1 / INRF2 is a key regulator of the NRF2 transcription factor, which transactivates the antioxidant response element (ARE) and upregulates numerous proteins involved in antioxidant defense. Under basal conditions, KEAP1 / INRF2 targets NRF2 for ubiquitination and proteolytic degradation and as such is responsible for the rapid turnover of NRF2. KEAP1 / INRF2 retains NFE2L2 / NRF2 in the cytosol. KEAP1 / INRF2 functions as substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1. It targets NFE2L2 / NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. KEAP1 / INRF2 may also retain BPTF in the cytosol. It targets PGAM5 for ubiquitination and degradation by the proteasome.

配方: PBS

分子量: 69.7 kDa

序列: Gln2-Cys624

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.