
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Rat CADM3 / NECL1 / IGSF4B Protein (Fc tag)

产品货号: TP08224

第三版

别名:CADM3;Igsf4b;Necl1

描述:Cell Adhesion Molecules (CAMs) are proteins located on the cell surface involved with the binding with other cells or with the extracellular matrix (ECM) in the process called cell adhesion. These proteins are typically transmembrane receptors and are composed of three domains: an intracellular domain that interacts with the cytoskeleton, a transmembrane domain, and an extracellular domain that interacts either with other CAMs of the same kind (homophilic binding) or with other CAMs or the extracellular matrix (heterophilic binding). Cell adhesion molecule 3, also known as Immunoglobulin superfamily member 4B, CADM3, and NECL1, is a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule which has Ca(2+)-independent homo- or heterophilic cell-cell adhesion activity and plays an important role in the formation of synapses, axon bundles and myelinated axons. Isoform 1 of CADM3 is expressed mainly in adult and fetal brain. Isoform 2 of CADM3 is highly expressed in adult brain and weakly expressed in placenta. In brain, Isoform 2 is highly expressed in cerebellum. CADM3 is involved in the cell-cell adhesion. It has both calcium-independent homophilic cell-cell adhesion activity and calcium-independent heterophilic cell-cell adhesion activity with IGSF4, PVRL1 and PVRL3. The interaction with EPB41L1 may regulate structure or function of cell-cell junctions. CADM3 may act as a tumor suppressor in glioma and loss of it in glioma may be caused by histone deacetylation.

配方:PBS

分子量:60.5 kDa

序列:Met1-His328

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C 或者-70 ° C. 避免反复冻融.