

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Recombinant Human CSF1R / MCSF Receptor / CD115 Protein (His & Fc tag)

产品货号: TP08386

第三版

别名:C-FMS;CD115;CSF-1R;CSFR;FIM2;FMS;HDLS;M-CSF-R;MCSF Receptor

描述:M-CSFR encoded by the proto-oncogene c-fms is the receptor for colony stimulating factor 1 (CSF1R), a cytokine involved in the proliferation, differentiation, and activation of macrophages. This cell surface glycoprotein is consisted by an extracellular ligand-binding domain, a single membrane-spanning segment, and an intracellular tyrosine kinase domain. Binding of CSF1 activates the receptor kinase, leading to "autophosphorylation" of receptor subunits and the concomitant phosphorylation of a series of cellular proteins on tyrosine residues. CSF1R is a tyrosine kinase receptor that is absolutely required for macrophage differentiation and thus occupies a central role in hematopoiesis. CSF1 and its receptor (CSF1R, product of c-fms proto-oncogene) were initially implicated as essential for normal monocyte development as well as for trophoblastic implantation. This apparent role for CSF1/CSF1R in normal mammary gland development is very intriguing because this receptor correlates with tumor cell invasiveness and adverse clinical prognosis. Tumor cell expression of CSF1 and its receptor correlates with tumor cell invasiveness and adverse clinical prognosis. Tumor cell expression of CSF1R is under the control of several steroid hormones (glucocorticoids and progestins) and the binding of several bHLH transcription factors, while tumor cell expression of CSF1 appears to be regulated by other hormones, some of which are involved in normal lactogenic differentiation. However, studies have demonstrated that CSF1 and CSF1R have additional roles in mammary gland development that may elucidate potential relationships of growth factor-induced biological changes in the breast during pregnancy and tumor progression.

配方:PBS

分子量:82.4 kDa

序列:Met 1-Glu512

纯度:> 95% by HPLC

浓度:

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4°C保存 (1-2周). 长期保存在-20°C或者-70°C. 避免反复冻融.