

---

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

---

Recombinant Human TPL2 / MAP3K8 / MEKK8 Protein (GST tag)

产品货号: TP08403

---

第三版

别名:AURA2;c-COT;COT;EST;ESTF;MEKK8;Tpl-2;TPL2

**描述:** Mitogen-activated protein kinase kinase kinase 8, also known as Cancer Osaka thyroid oncogene, Proto-oncogene c-Cot, Serine/threonine-protein kinase cot, Tumor progression locus 2 and MAP3K8, is a cytoplasm protein which belongs to the protein kinase superfamily, STE Ser/Thr protein kinase family and MAP kinase kinase kinase subfamily. MAP3K8 is expressed in several normal tissues and human tumor-derived cell lines. Isoform 1 of MAP3K8 is activated specifically during the S and G2/M phases of the cell cycle. MAP3K8 is required for TLR4 activation of the MEK/ERK pathway. It is able to activate NF-kappa-B 1 by stimulating proteasome-mediated proteolysis of NF-kappa-B 1/p105. MAP3K8 plays a role in the cell cycle. The longer form has some transforming activity, although it is much weaker than the activated cot oncoprotein. MAP3K8 oncogene linked to human endometrial carcinoma suggesting that it may be another molecule involved in human endometrial cancer. MAP3K8 may also be an important mediator of intracellular mechanotransduction in human bone marrow-derived mesenchymal stem cells (MSCs).

**配方:** PBS

**分子量:** 68 kDa

**序列:** Met 30-Arg 397

**纯度:** > 95% by HPLC

**浓度:**

**内毒素:** <1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.