
本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Recombinant Human IL-1RAcP / IL-1R3 Protein (His & Fc tag)

产品货号: TP08427

第三版

别名:C3orf13;IL-1RAcP;IL1R3

描述: Interleukin-1 receptor accessory protein (IL-1RAcP) also known as Interleukin-1 receptor member 3 (IL-1R3) is a cytokine receptor which binds interleukin 1. The IL-1 receptor accessory protein (IL1RAP) is a transmembrane protein that interacts with IL-1R and is required for IL-1 signal transduction. Interleukin 1 induces synthesis of acute phase and proinflammatory proteins during infection, tissue damage, or stress, by forming a complex at the cell membrane with an interleukin 1 receptor and an accessory protein. IL-1RAcP/IL-1R3 is a necessary part of the interleukin 1 receptor complex which initiates signalling events that result in the activation of interleukin 1-responsive genes. Alternative splicing of this gene results in two transcript variants encoding two different isoforms, one membrane-bound and one soluble. The ratio of soluble to membrane-bound forms increases during acute-phase induction or stress. IL-1RAcP/IL-1R3 mediates interleukin-1-dependent activation of NF-kappa-B. Isoform 1 is part of the membrane-bound form of the IL-1 receptor. Signaling involves formation of a ternary complex containing IL1R1, TOLLIP, MYD88, and IRAK1 or IRAK2. Isoform 2 modulates the response to interleukins by associating with soluble IL1R1 and enhancing interleukin-binding to the decoy receptor.

配方: PBS

分子量: 67.3 kDa

序列: Met 1-Glu 359

纯度: > 95% by HPLC

浓度:

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.