

---

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

---

Recombinant Mouse TRAIL R2 / CD262 / TNFRSF10B Protein (His & Fc tag)

产品货号: TP08492

---

第三版

别名:DR5;KILLER;Ly98;MK;TRAILR2;TRICK2A;TRICK2B;TRICKB

**描述:** Tumor necrosis factor receptor superfamily, member 10b, official symbol TNFRSF10B, also known as Death receptor 5, CD262, TNF-related apoptosis-inducing ligand receptor 2 (TRAIL R2), is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. TRAIL R2/CD262/TNFRSF10B was purified independently as the only receptor for TRAIL detectable on the surface of two different human cell lines that undergo apoptosis upon stimulation with TRAIL. TRAIL R2/CD262/TNFRSF10B contains two extracellular cysteine-rich repeats, typical for TNF receptor (TNFR) family members, and a cytoplasmic death domain. TRAIL R2/CD262/TNFRSF10B mediates apoptosis via the intracellular adaptor molecule FADD/MORT1. TRAIL receptors can signal both death and gene transcription, functions reminiscent of those of TNFR1 and TRAMP, two other members of the death receptor family. Defects in TRAIL R2/CD262/TNFRSF10B may be a cause of head and neck squamous cell carcinomas (HNSCC) also known as squamous cell carcinoma of the head and neck.

**配方:** PBS

**分子量:** 41.8 kDa

**序列:** Met 1-Ser 177

**纯度:** > 95% by HPLC

**浓度:**

**内毒素:** <1.0 EU per 1 ug of protein (determined by LAL method)

**存储:** +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.