

Anti-Human/Monkey Cleaved-NOTCH4 (V1432) Polyclonal Antibody**Polyclonal Antibody****Cat.NO.: PA02158**

3th Edition

Description: This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In *Drosophila*, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network, and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands, and may play a role in vascular, renal and hepatic development. This gene may be associated with susceptibility to schizophrenia in a small portion of cases. An alternative splice variant has been described but its biological nature has not been determined.

Antigen: Synthesized peptide derived from the C-terminal region of human Notch 4

Form:

How to use: 1.0 ml distilled water will be added to the product

Stability: Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

Dilution: PBS (pH7.4) containing 1% BSA

Application: This antibody can be used for western blotting in concentration of 1?5?g/ml.

Specificity: Highly expressed in the heart, moderately in the lung and placenta and at low levels in the liver, skeletal muscle, kidney, pancreas, spleen, lymph node, thymus, bone marrow and fetal liver. No expression was seen in adult brain or peripheral blood leukocytes.