

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Anti-Human CSE1L Polyclonal Antibody

**Polyclonal Antibody** 

Cat.NO.: PA02443

3th Edition

**Description:** Proteins that carry a nuclear localization signal (NLS) are transported into the nucleus by the importinal pha/beta heterodimer. Importinal pha binds the NLS, while importinate mediates translocation through the nuclear pore complex. After translocation, RanGTP binds importinate and displaces importinal pha. Importinal pha must then be returned to the cytoplasm, leaving the NLS protein behind. The protein encoded by this gene binds strongly to NLS-free importinal pha, and this binding is released in the cytoplasm by the combined action of RANBP1 and RANGAP1. In addition, the encoded protein may play a role both in apoptosis and in cell proliferation. Alternatively spliced transcript variants have been found for this gene.

Antigen: Synthesized peptide derived from the N-terminal region of human CAS

Form:

How to use:1.0 ml distilled water will be added to the product

**Stability:** Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

Dilution: PBS (pH7.4) containing 1% BSA

Application: This antibody can be used for western blotting in concentration of 1?5?g/ml.

**Specificity:**Highly expressed in proliferating cells.

1/1