

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Anti-Human/Mouse/Rat Phospho-BCL-x (Thr47) Polyclonal Antibody

Polyclonal Antibody

Cat.NO.: PA04063

3th Edition

Description:BCL2L1 is a member of he BCL-2 protein family. BCL2L1 is expressed as two distinct isoforms, Bcl-xL and Bcl-xS, and is located at the outer mitochondrial membrane. The Bcl-XL isoform is a 233 amino acid protein, acting as an apoptotic inhibitor. Bcl-XL can forms heterodimers with BAX, BAK or BCL2, and the heterodimerization with BAX does not seem to be required for anti-apoptotic activity. The Bcl-XS isoform is a shorter variant that is 178 amino acids in length and lacks a 63 amino acid region (amino acids 126-188), acting as an apoptotic activator. This antibody can recognize Bcl-XL, Bcl-XS and Bcl-X(Beta).

Antigen: Synthesized peptide derived from human Bcl-x around the phosphorylation site of T47.

Form:

How to use:1.0 ml distilled water will be added to the product

Stability: Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

Dilution: PBS (pH7.4) containing 1% BSA

Application: This antibody can be used for western blotting in concentration of 1?5?g/ml.

Specificity:Bcl-X(S) is expressed at high levels in cells that undergo a high rate of turnover, such as developing lymphocytes. In contrast, Bcl-X(L) is found in tissues containing long-lived postmitotic cells, such as adult brain.

1/1