

Anti-Human/Mouse/Rat JAK2 Polyclonal Antibody**Polyclonal Antibody****Cat.NO.: PA04570**

3th Edition

Description:Non-receptor tyrosine kinase involved in various processes such as cell cycle progression, apoptosis, mitotic recombination, genetic instability and histone modifications. In the cytoplasm, plays a pivotal role in signal transduction via its association with cytokine receptors, which constitutes an initiating step in signaling for many members of the cytokine receptor superfamily including the receptors for growth hormone (GHR), prolactin (PRLR), leptin (LEPR), erythropoietin (EPOR), granulocyte-macrophage colony-stimulating factor (CSF2), thrombopoietin (THPO) and multiple interleukins. Following stimulation with erythropoietin (EPO) during erythropoiesis, it is autophosphorylated and activated, leading to its association with erythropoietin receptor (EPOR) and tyrosine phosphorylation of residues in the EPOR cytoplasmic domain. Also involved in promoting the localization of EPOR to the plasma membrane. Also acts downstream of some G-protein coupled receptors. Plays a role in the control of body weight (By similarity). Mediates angiotensin-2-induced ARHGEF1 phosphorylation. In the nucleus, plays a key role in chromatin by specifically mediating phosphorylation of 'Tyr-41' of histone H3 (H3Y41ph), a specific tag that promotes exclusion of CBX5 (HP1 alpha) from chromatin.

Antigen:Recombinant Protein of JAK2

Form:

How to use:1.0 ml distilled water will be added to the product

Stability: Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

Dilution:PBS (pH7.4) containing 1% BSA

Application:This antibody can be used for western blotting in concentration of 1?5?g/ml.

Specificity:Expressed in blood, bone marrow and lymph node.