

**Anti-Human CCL15 Polyclonal Antibody**

**Polyclonal Antibody**

**Cat.NO.: PA04877**

---

3th Edition

**Description:** This gene, chemokine (C-C motif) ligand 15, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene is chemotactic for T cells and monocytes and induces N-acetyl-beta-D-glucosaminidase release in monocytes. It induces changes in intracellular calcium concentration in monocytes and is thought to act through the CCR1 receptor. Read-through transcripts are expressed that include exons from the downstream cytokine gene, chemokine (C-C motif) ligand 14, and are represented as GeneID: 348249.

**Antigen:** Synthetic peptide from human protein

**Form:**

**How to use:** 1.0 ml distilled water will be added to the product

**Stability:** Lyophilized product, 5 years at 2 – 8°C; Solution, 2 years at –20°C

**Dilution:** PBS (pH7.4) containing 1% BSA

**Application:** This antibody can be used for western blotting in concentration of 1?5?g/ml.

**Specificity:** Most abundant in heart, skeletal muscle and adrenal gland. Lower levels in placenta, liver, pancreas and bone marrow. CCL15(22-92), CCL15(25-92) and CCL15(29-92) are found in high levels in synovial fluids from rheumatoid patients.