

**CXCL17, 22-119aa Human, His tag, E.coli**

**Cat.NO.: TP01797**

3th Edition

**Synonyms:** Chemokine (C-X-C motif) ligand 17 , Dcip1, DMC, MGC138300, UNQ473, VCC-1, VCC1

**Description:** Chemokine (C-X-C motif) ligand 17, also known as CXCL17, is a small cytokine belonging to the CXC chemokine family. CXCL17 is a 119 amino acid secreted protein that plays a role in angiogenesis. CXCL17 is expressed in skeletal muscle, trachea, lung, intestine and stomach, and is upregulated in duodenal mucosa of patients with acute cholera, as well as breast tumors. CXCL17 is considered a housekeeping chemokine for the movement of immature dendritic cells and non activated blood monocytes into tissues, and is thought to be involved in the innate immune response. Recombinant human CXCL17 protein, fused to His-tag at N-terminus, was expressed in E.coli.

**Form:** Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

**Molecular Weight:** 13.7 kDa (119aa)

**Sequences:**

MGSSHHHHHHSSGLVPRGSHMSSLNPGVARGHRDRGQASRRWLQEGGQECECKDWFLRAPRRKFMTVSGLPK  
KQCPCDHFKGNVKKTRHQRHHRKPNKHSRACQQFLKQCQLRSFALPL

**Purity:** > 95% by HPLC

**Concentration:** 0.5 mg/ml (determined by Bradford assay)

**Endotoxin Level:** <1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:** Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.